

Each drug is recommended for use in combination with other antiretrovirals (except possibly Atripla and rilpivirine/Truvada). Dose may need to be adjusted; see drug page or package insert for complete information. Drugs shown are actual size, except for Fuzeon. Drug chart information is current as of Aug. 1, 2011. Thanks to Swarup Mehta, PharmD, Walgreens pharmacist. For a list of Walgreens pharmacies that specialize in HIV/AIDS care and other resources, visit www.HIV.walgreens.com/chart. Using a QR code reader app, you can also scan the QR code at right with your iPhone or other smartphone to go directly to the website. See the POSITIVELY AWARE 2011 HIV Drug Guide at www.positivelyaware.com for more information.



SPONSORED BY
Walgreens

NUCLEOSIDE/NUCLEOTIDE ANALOG REVERSE TRANSCRIPTASE INHIBITOR (NRTI, NNRTI)



Combivir

lamivudine/zidovudine, or 3TC/AZT

ADULT SINGLE DOSE: One tablet (150 mg lamivudine, 300 mg zidovudine), twice daily. No food restrictions (can be taken with or without food).



Epivir

lamivudine, or 3TC

ADULT SINGLE DOSE: One 300 mg tablet, once daily; or one 150 mg tablet, twice daily. No food restrictions (can be taken with or without food).



Retrovir

zidovudine, AZT, or ZDV

ADULT SINGLE DOSE: One 300 mg tablet, twice daily. No food restrictions (can be taken with or without food).



Truvada

tenofovir DF/emtricitabine, or TDF/FTC

ADULT SINGLE DOSE: One tablet (300 mg tenofovir DF, 200 mg emtricitabine), once daily. No food restrictions (can be taken with or without food).



Viread

tenofovir disoproxil fumarate, or TDF

ADULT SINGLE DOSE: One 300 mg tablet, once daily. No food restrictions (can be taken with or without food).



Ziagen

abacavir sulfate, or ABC

ADULT SINGLE DOSE: Two 300 mg tablets, once daily; or one 300 mg tablet, twice daily. No food restrictions (can be taken with or without food).



Intelence

etravirine, or ETR

ADULT SINGLE DOSE: One 200 mg tablet, twice daily; or two 100 mg tablets, twice daily. Take with food.



Sustiva

efavirenz, or EFV

ADULT SINGLE DOSE: One 600 mg tablet, once daily; or three 200 mg capsules, once daily, preferably at bedtime. Take on an empty stomach, or may be taken with a light, low-fat snack.



Atripla

efavirenz/tenofovir DF/emtricitabine, or EFV/TDF/FTC

ADULT SINGLE DOSE: One tablet (600 mg efavirenz, 300 mg tenofovir DF, 200 mg emtricitabine), once daily, preferably at bedtime. Take on an empty stomach, or may be taken with a light, low-fat snack.



Aptivus

tipranavir, or TPV

ADULT SINGLE DOSE: Two 250 mg capsules with two 100 mg tablets Norvir, twice daily. Aptivus must be dosed with Norvir. No food restrictions (can be taken with or without food); take with food to improve Norvir tolerability.



Invirase

saquinavir, or SQV

ADULT SINGLE DOSE: Two 500 mg tablets with one 100 mg tablet Norvir, twice daily. Invirase must be dosed with Norvir. Take with food, or within two hours after a meal.



Lexiva

fosamprenavir calcium, or FPV

ADULT SINGLE DOSE: Two 700 mg tablets with 100 mg or 200 mg Norvir, once daily; or two 700 mg tablets without Norvir, twice daily; if protease inhibitor experienced, 700 mg with 100 mg Norvir, twice daily. No food restrictions (can be taken with or without food).



Prezista

darunavir, or DRV


ADULT SINGLE DOSE: Two 400 mg tablets, once daily for first-time therapy and treatment-experienced people with no Prezista-associated drug resistance; or one 600 mg tablet, twice daily for treatment-experienced people with at least one Prezista-associated drug mutation. All doses taken with 100 mg Norvir. Take with food.



Viracept

nelfinavir, or NFV

ADULT SINGLE DOSE: Two 625 mg tablets, twice daily; or five 250 mg tablets, twice daily. Take with food, preferably high in fat.



Fuzeon

enfuvirtide, T-20, or ENF

ADULT SINGLE DOSE: 90 mg subcutaneous injection, twice daily. No food restrictions (can be taken with or without food).

NUCLEOSIDE/NUCLEOTIDE ANALOG REVERSE TRANSCRIPTASE INHIBITOR (NRTI, NNRTI)



Emtriva

emtricitabine, or FTC

ADULT SINGLE DOSE: One 200 mg capsule, once daily. No food restrictions (can be taken with or without food).



Epzicom

abacavir/lamivudine, or ABC/3TC

ADULT SINGLE DOSE: One tablet (600 mg abacavir, 300 mg lamivudine), once daily. No food restrictions (can be taken with or without food).



Trizivir

abacavir/lamivudine/zidovudine, or ABC/3TC/AZT

ADULT SINGLE DOSE: One tablet (300 mg abacavir, 150 mg lamivudine, 300 mg zidovudine), twice daily. No food restrictions (can be taken with or without food).



Videx EC

didanosine, or ddl

ADULT SINGLE DOSE: One 400 mg capsule, once daily with adjustments for weight and drug interactions (see the Drug Guide). Take on an empty stomach, 30 minutes before or two hours after food or drink, except water. (Generic pictured.)



Zerit

stavudine, or d4T

ADULT SINGLE DOSE: One 40 mg capsule, twice daily with adjustments for weight (see the Drug Guide). No food restrictions (can be taken with or without food).



Edurant

rilpivirine, or RPV

ADULT SINGLE DOSE: One 25 mg tablet, once daily. Take with food (at least 500 calories).



Rescriptor

delavirdine, or DLV

ADULT SINGLE DOSE: Two 200 mg tablets, three times a day; or four 100 mg tablets, three times a day. No food restrictions (can be taken with or without food).



Viramune

nevirapine, or NVP

ADULT SINGLE DOSE: One 200 mg immediate-release (IR) tablet, once daily for the first 14 days. Then, one 400 mg extended-release (XR) tablet, once daily; or one 200 mg IR tablet, twice daily; or two 200 mg IR tablets, once daily. No food restrictions (can be taken with or without food).

rilpivirine/Truvada

rilpivirine/tenofovir DF/emtricitabine

ADULT SINGLE DOSE: One tablet combination (25 mg rilpivirine and Truvada [300 mg tenofovir DF, 200 mg emtricitabine]), once daily has been selected in the Phase 3 trials. Take with food (at least 500 calories). (Not yet approved at press time; photo unavailable.)

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR (NNRTI)

FIXED DOSE REGIMEN

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR (NNRTI)

FIXED DOSE REGIMEN

PROTEASE INHIBITOR (PI)

ENTRY INHIBITOR

PROTEASE INHIBITOR (PI)

INTEGRASE INHIBITOR

ENTRY INHIBITOR



Crixivan

indinavir, or IDV

ADULT SINGLE DOSE: Two 400 mg capsules, every 8 hours; or two 400 mg capsules with 100 mg Norvir twice a day. Take with no food or a low-fat snack (with food with Norvir); drink at least 48 oz. of water daily.



Kaletra

lopinavir/ritonavir, or LPV/r

ADULT SINGLE DOSE: Four tablets (200 mg lopinavir/50 mg ritonavir), once daily if fewer than 3 lopinavir resistance mutations; or two 200/50 mg tablets, twice daily. Half-strength film-coated tablet available: 100 mg lopinavir/25 mg ritonavir. No food restrictions; take with food to improve Norvir tolerability.



Norvir

ritonavir, or RTV

ADULT SINGLE DOSE: Used mostly for boosting, 100 mg-400 mg once or twice daily with another PI. Must be taken with food.



Reyataz

atazanavir sulfate, or ATV

ADULT SINGLE DOSE: One 300 mg capsule plus 100 mg Norvir, once daily or two 200 mg capsules, once daily (see Drug Guide). Take with food.



Isentress

raltegravir, or RAL

ADULT SINGLE DOSE: One 400 mg tablet, twice daily. No food restrictions (can be taken with or without food).



Selzentry

maraviroc, or MVC

ADULT SINGLE DOSE: 150, 300, or 600 mg (available in 150 and 300 mg tablets), twice daily; dose differs based upon other HIV medications used (see Drug Guide). No food restrictions (can be taken with or without food).